

## The Importance of DNA methylation in Prostate Cancer development

Massie, C. E., Mills, I. G., & Lynch, A. G. (2016). The Importance of DNA methylation in Prostate Cancer development. *Journal of Steroid Biochemistry and Molecular Biology*.  
<https://doi.org/10.1016/j.jsbmb.2016.04.009>

**Published in:**

Journal of Steroid Biochemistry and Molecular Biology

**Document Version:**

Publisher's PDF, also known as Version of record

**Queen's University Belfast - Research Portal:**

[Link to publication record in Queen's University Belfast Research Portal](#)

**Publisher rights**

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution and reproduction in any medium, provided the author and source are cited.

**General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).



Contents lists available at ScienceDirect

# Journal of Steroid Biochemistry and Molecular Biology

journal homepage: [www.elsevier.com/locate/jsbmb](http://www.elsevier.com/locate/jsbmb)



## Review

# The importance of DNA methylation in prostate cancer development

Charles E. Massie<sup>a</sup>, Ian G. Mills<sup>b,c,d</sup>, Andy G. Lynch<sup>a,\*</sup>

<sup>a</sup> Cancer Research UK Cambridge Institute, University of Cambridge, Li Ka Shing Centre, Robinson Way, Cambridge, UK

<sup>b</sup> Prostate Cancer Research Group, Centre for Molecular Medicine (Norway), University of Oslo and Oslo University Hospitals, Gaustadalleen, Oslo, Norway

<sup>c</sup> Department of Molecular Oncology, Oslo University Hospitals, Oslo, Norway

<sup>d</sup> PCUK/Movember Centre of Excellence for Prostate Cancer Research, Centre for Cancer Research and Cell Biology (CCRCB), Queen's University Belfast, Belfast, UK

## ARTICLE INFO

### Article history:

Received 6 February 2016

Received in revised form 9 April 2016

Accepted 17 April 2016

Available online xxx

### Keywords:

Prostate cancer

Epigenetic

Methylation

Biomarkers

Metabolism

Mitochondria

## ABSTRACT

After briefly reviewing the nature of DNA methylation, its general role in cancer and the tools available to interrogate it, we consider the literature surrounding DNA methylation as relating to prostate cancer. Specific consideration is given to recurrent alterations. A list of frequently reported genes is synthesized from 17 studies that have reported on methylation changes in malignant prostate tissue, and we chart the timing of those changes in the diseases history through amalgamation of several previously published data sets.

We also review associations with genetic alterations and hormone signalling, before the practicalities of investigating prostate cancer methylation using cell lines are assessed. We conclude by outlining the interplay between DNA methylation and prostate cancer metabolism and their regulation by androgen receptor, with a specific discussion of the mitochondria and their associations with DNA methylation.

© 2016 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

## Contents

1. Introduction: why consider the epigenome?.....	00
1.1. The origins of prostate cancer are not obviously genomic .....	00
1.2. Considerations for DNA methylation profiling in prostate cancer .....	00
1.3. The data used in this review .....	00
2. Recurrent epigenetic changes in prostate cancer: markers and drivers of disease evolution .....	00
2.1. Early epigenetic changes in prostate carcinogenesis .....	00
2.2. An epigenetic 'field-effect' in cancerous prostates .....	00
2.3. Underlying clonal stability and ongoing epigenetic evolution in prostate tumours .....	00
2.4. DNA methylation markers for the detection and stratification of prostate cancer .....	00
2.5. Epigenetic regulation of alternative promoter usage in prostate cancer .....	00
2.6. Associations between epigenetic and genetic alterations in prostate cancer .....	00
2.7. Epigenetic changes impacting on hormone signalling in prostate cancer .....	00
2.8. Prostate cancer cell lines .....	00
3. Effects of the metabolic pool on DNA methylation.....	00
4. Mitochondrial methylation and prostate cancer .....	00
4.1. Importance of mitochondria in PCa .....	00
4.2. Reciprocal regulation of nuclear methylation and mitochondria .....	00
5. Current perspectives and future directions for the role of epigenomic changes in prostate cancer .....	00
Acknowledgements.....	00
Appendix A. Supplementary data .....	00
References .....	00

\* Corresponding author.

E-mail address: [andy.lynch@cruk.cam.ac.uk](mailto:andy.lynch@cruk.cam.ac.uk) (A.G. Lynch).

## 1. Introduction: why consider the epigenome?

### 1.1. The origins of prostate cancer are not obviously genomic

Cancer genomics studies have identified recurrently mutated genes and mutation hotspots in a number of cancer types. However, such studies in prostate adenocarcinomas have identified no genes recurrently mutated in more than a seventh of cases [1]. Studies of locally advanced and metastatic prostate cancer have revealed extensive intratumoural clonal heterogeneity [2–6], in some cases revealing clones with distinct genomic origins [3,5]. This extensive clonal and spatial heterogeneity creates a significant sampling problem for studies that rely on the use of single tumour specimens. In such studies, intratumoural heterogeneity will amplify intertumoural heterogeneity, contributing to the low recurrence rates of genes affected by point mutations in prostate cancer [1].

Other mutation types have higher levels of recurrence, e.g. 8p deletions (in 40% of cases) and *TMPRSS2-ERG* fusions (in 50% of cases) [7,8], suggesting that these may be early or convergent events in prostate tumourigenesis. However, many prostate tumours have no definitive genomic driver event [1]. This is consistent with the existence of events that precede the first somatic point mutations and possibly also the acquisition of the first copy number and structural variants. Epigenomic changes are among the candidates for early events. Of these, DNA methylation changes have been widely studied and found to be the most recurrent events in both locally advanced and metastatic prostate tumours [9,10].

In prostate cancer, recurrent genome-wide and locus specific DNA methylation alterations have been known for decades [11,12] and these events impact on gene expression potential [12,13]. The high recurrence rates of specific somatic alterations in DNA methylation support a strong selective pressure for these events and implicate them in the development of neoplastic phenotypes and as rate limiting steps in disease evolution [12,13]. Cytosine methylation is the most widely studied epigenetic marker in cancer due to the development of quantitative genomics methods that are compatible with tissue samples obtained from surgical specimens. To date most studies have used prostate cancer cell lines when profiling chromatin structures and histone modifications [14–16] and other variants of cytosine modification have only been assessed at a global level in primary prostate cancer tissue [17]. Therefore for the purposes of this review we will restrict our focus to the wealth of studies that have profiled cytosine methylation in prostate cancer.

### 1.2. Considerations for DNA methylation profiling in prostate cancer

DNA methylation is a stable, heritable genome modification that can provide insights into a tumour's origins and evolution. Methylation profiling is aided by the number of well-developed techniques and analysis methods available. It is aided also by the requirement only for standard preparation of genomic DNA as input, making it applicable to routine tumour tissue collections (i.e. in contrast to methods that require cross-linked chromatin).

A range of methylation profiling methods have been developed, from cis-linked, base-pair resolution bisulfite sequencing of the whole genome (WGBS [18]) or GC-base enriched regions (eRRBS [19,20]), to array based averaging of methylation at specific CpG sites [21,22], to locus-averaging methods that identify methylation 'peaks' (me-DIP [23,24]) or that may be a proxy for functional methylation changes (e.g. MBD pull-down [25,26]). The most widely used platform for studies of clinical tissue samples is the Infinium 450k array [21], that continues to be used due to its reproducibility, well developed analysis methods and consequent potential for

integration with a wealth of published data from this platform [27,28].


Sequencing methods provide the highest resolution profiles and cis-linkage information about the status of adjacent CpGs on the same strand, but have the largest analytical burden. Standard bisulfite sequencing methods do not, however, discriminate between methyl-cytosine (5-mC) and hydroxy-methyl-cytosine (5-hmC), although these marks are believed to have different functional consequences. To discriminate 5-hmC and 5-mC a two stage analysis is required comparing the results of bisulfite sequencing (for a combined 5-hmC and 5-mC signal) and oxidative bisulfite sequencing (for 5-hmC alone), followed by subtractive analysis. Alternatively, me-DIP approaches using 5-hmC and 5-mC specific antibodies can also discriminate these signals to provide locus-averaged signals. Future improvements in sequencing yields and sensitivities for single molecule sequencing platforms may provide different insights into the epigenetic landscape, for example long read technologies may allow better phasing of epigenetic states along chromosome domains. Recent reports suggest that nanopore-based sequencing technologies may be able directly to read the 5-mc or 5-hmc modifications of cytosine [29].

The DNA methylation landscape varies across the genome, generally showing higher methylation at repeat sequences and retrotransposons compared to lower methylation at active gene promoters and CpG-islands [18,30]. High levels of DNA methylation at gene promoter regions (and around the transcription start site) correlates with low gene expression [12,18]. Within a gene locus the methylation profile can vary widely (Fig. 1A), meaning that comparisons between samples (i.e. differential analysis) must rely on either comparisons of individual CpG sites or by defining local methylation domains (e.g. differentially methylated regions, DMRs) [31]. CpG-islands have low DNA methylation variance in cancer, while adjacent regions (termed CpG-shelves and CpG-shores; Fig. 1B) tend to show higher variation. Most recently locally disordered methylation or epipolymorphisms have been reported [32,33] and linked to evolutionary plasticity in cancer, as previously suggested for epigenetic variation [33–35].

The selection of samples for cancer genome sequencing is usually simple because the aim is to identify somatically acquired changes (e.g. comparing tumour tissue with a germline control sample – often blood or buccal swabs). However, tissue specific methylation profiles mean that the most appropriate control sample for cancer methylome studies is normal tissue from the same organ. More stringently, one might aim to match the proportions of cell types (e.g. epithelial, stromal, immune) in the tumour and normal tissue samples. In many cancer types (including prostate cancer) a 'field-effect' change has been observed in the tumour adjacent normal tissue, consistent with a pre-neoplastic state. Therefore, depending on the study aims it may be most appropriate to compare epigenetic state between tumours, tumour adjacent normal tissue and age-matched tumour-free normal tissue.

### 1.3. The data used in this review

Through this review we will illustrate key points using previously published data sets. For ease of comparison, and due to their greater number, we will focus solely on data generated using the Illumina Infinium HumanMethylation450 BeadChip. For individual genes, and to relate methylation levels to gene expression we will use the 'TCGA' prostate adenocarcinoma data [28], interrogated and plotted using the TCGA Wanderer interface [36]. For consistency we use Wanderer's associations of probes to genes throughout, although this naturally leads to probes mapping to multiple genes.



**Fig. 1.** (A) Average methylation profiles for prostate tumours ( $n = 340$ ) and normal prostate tissue ( $n = 49$ ) at the *GSTP1* gene locus using the TCGA data set (see Section 1.3). (B) Schematic showing the *GSTP1* gene locus, indicating the location of the CpG-island, CpG-shore (<2kb from island) and CpG-shelf (2–4kb from island). (C) Correlation scatter plot for *GSTP1* expression and methylation (using the 450k array probe highlighted in panel-A). (D) Correlation scatter plot for *CDKN2A* expression and methylation (using the 450k array probe highlighted in panel-E). (E) Average methylation profiles for prostate at the *CDKN2A* gene locus using the Prostate TCGA data set (see Section 1.3). (F) Schematic showing the gene and CPGI features at the *CDKN2A* locus.

For the second data set, the ‘Tissue’ data set, we amalgamate data from several sources [26,27,37–39] to obtain methylation statuses for prostates from men with no prostate cancer (“Normal”), morphologically normal tissue from men with prostate cancer (“Benign”), benign prostatic hyperplasia (“Hyperplasia”), neoplastic tissue (“Neoplasia”), primary tumours (“Tumour”) and

metastases (“Metastasis”). We also obtain blood profiles [39] as an additional reference.

For Fig. 3A, where space is a constraint, we use only a subset of these drawn from two sources [26,37]. The third data set, the ‘Cell line’ data set consists of the combined HumanMethylation450 data detailed later in Table 1. Finally, to annotate genes with

**Table 1**  
Public prostate cell line methylation data (January 2016).

	GoldenGate	27k	450k	Nimblegen	MBD-seq	RRBS	NOMe-seq	MRE-seq	MEDIP
DU-145	GSM1125684, GSM696035	GSM573668	GSM1633638, GSM1323599		SRX118022, SRX118022, GSM1050087, GSM1050090, GSM1050093, GSM1050096, GSM1050099	GSM1050102			
PC3	GSM1125685, GSM696036	GSM573670	GSM1633598, GSM1519011 to GSM1519016*, GSM1323600	GSM1142996 to GSM1143004	GSM1050088, GSM1050091, GSM1050094, GSM1050097, GSM1050100	GSM1050103	GSM1383852		
LNCaP	GSM1125683, GSM696034	GSM573669 (FGC)	GSM1519017, GSM1519018, GSM999368, GSM847569 to GSM847571	GSM1142987 to GSM1142995*	SRX118021, GSM605080, GSM605081	GSM683768, GSM683776, GSM683862, GSM683863, GSM683924, GSM683946		GSM684592, GSM684597 to GSM684600	GSM605948, GSM605950, GSM605954 to GSM605956
ARCaP	GSE35246 (27 samples*)								
PrEC	GSE35246 (4 samples*)		GSM999369, GSM847572 to GSM847574	GSM1142978 to GSM1142986*	GSM605082, GSM605083	GSM683760, GSM683838	GSM1383851	GSM684593 to GSM684596	GSM605945 to GSM605947, GSM605951 to GSM605953
PrED					SRX118020				
RWPE-1		GSM573671	GSM1323601						

Detailing the data available for common prostate cell lines from Gene Express Omnibus (GSM/GSE) and the Short Read Archive (SRX). \* – includes control and treated cell lines. Platforms included include Illumina Goldengate Methylation Cancer Panel I (“GoldenGate”), Illumina Infinium HumanMethylation27 BeadChip (“27k”), Illumina Infinium HumanMethylation450 BeadChip (“450k”), Nimblegen Human DNA Methylation 3x720K CpG Island Plus RefSeq promoter array (“Nimblegen”), methyl-CpG binding domain protein-enriched genome sequencing (“MBD-seq”), Reduced representation Bisulfite sequencing (“RRBS”), Nucleosome Occupancy and Methylome sequencing (“NOMe-seq”), Methylation-sensitive Restriction Enzyme Sequencing (“MRE-seq”), and Methylated DNA Immunoprecipitation Sequencing (“MEDIP”).



androgen receptor (AR) regulation data, we use two previously-published androgen-treated cell line time-course data sets [40,39].

## 2. Recurrent epigenetic changes in prostate cancer: markers and drivers of disease evolution

Recurrent alterations in DNA methylation at the *GSTP1* gene promoter and concomitant loss of *GSTP1* expression in prostate tumours were reported over 20 years ago [12]. This finding has been replicated in countless independent studies (for example [9,41,42]) and well over 1000 samples (reviewed in [43]), providing strong evidence that DNA methylation changes are indeed recurrent across patient cohorts and could be useful markers for the clinical detection of prostate cancer [9,44].

Several other genes have also been reported to be recurrently hypermethylated in prostate cancer by multiple studies. Synthesizing data from 17 studies [2,10,20,22,24,26–28,38,45–52], we identify 861 genes that are reported in two or more studies, 168 in three or more (detailed in Supplementary Table 1 and Supplementary Fig. 1), and 45 that are reported in four or more studies (Fig. 2). Some gene families are also recurrently affected, consistent with functional convergence, including multiple changes at the *HOX* gene family loci (Fig. 2 and Supplementary Fig. 1) [10,49].

### 2.1. Early epigenetic changes in prostate carcinogenesis

The high recurrence rates of these DNA methylation changes suggest that they may be early events in tumourigenesis. Indeed several studies have detected many such methylation changes in neoplastic samples (PIN) and tumour adjacent, morphologically benign tissue [53,54,27]. Indeed the majority of loci that have been suggested as differentiating benign and cancerous prostate appear already to have undergone epigenetic changes in neoplastic tissue (Fig. 2) impacting on their potential as prostate cancer markers, but highlighting early or shared events in cancer evolution.

Prospective multi-region sampling studies with good clinical annotations are therefore needed to map tumour specific markers comprehensively, in order to improve diagnostic accuracy from tissue biopsies and non-invasive monitoring. Given the current over-treatment of primary prostate cancer it is also imperative that robust panels of markers are developed to allow patient stratification for active surveillance or clinical intervention.

### 2.2. An epigenetic 'field-effect' in cancerous prostates

Recent studies suggest that DNA methylation changes in tumour adjacent 'normal' tissue may reflect a 'field effect' in cancerous prostates [50,24,27]. DNA methylation profiles have been reported to differ between tumour adjacent benign tissue and benign tissue from cancer-free prostates [50,24], while benign samples taken at different distances from prostate tumours show similar profiles, supporting a wide clonal expansion of morphologically normal cells [50].

This observation is consistent with the outgrowth of tumour clones originating from clonal benign and PIN tissue, a concept supported by a recent study comparing multiple benign, neoplasia and tumour samples from the same cancerous prostates [27]. In this study a common phylogenetic 'trunk' could be identified using either copy number or DNA methylation profiles, linking tumour samples, PIN and adjacent normal prostate samples. This indication of a shared clonal ancestry contrasts with the more sparse data from genome sequencing studies, where few point mutations, indels or structural variants link separate tumour foci and pre-cancerous tissues [5], suggesting that the expansion of genetically mutated

clones is a later event than the expansion of clones harbouring DNA methylation alterations.

Further studies are required to define more clearly the early neoplastic and tumour initiating events and also comprehensively to distinguish early events from convergent evolution. Longitudinal monitoring through the life-history of a patient with prostate cancer would be required to give a definitive answer to these questions, although this would be very difficult to achieve. An alternative approach would be to combine multi-region tissue sampling cohorts with base-pair resolution methylation sequencing to distinguish early events from convergent evolution. Understanding this would impact on the utility of these changes both as markers of early tumourigenesis and as targets for preventative medicine.

### 2.3. Underlying clonal stability and ongoing epigenetic evolution in prostate tumours

*GSTP1* methylation is present at all stages of prostate cancer development [51], showing that specific epigenetic changes can be maintained throughout disease evolution. In addition, methylome-wide studies have shown stable epigenetic profiles between metastatic deposits within a patient [26] and clear evidence of shared origins for metastatic deposits in primary tumours and premalignant lesions [27]. Interestingly, in most cases neoplastic lesions (PIN) were evolutionarily more similar to localized tumour samples, whereas metastatic deposits were often more closely related to a separate subset of localized tumour samples [27].


Hypomethylation of repeat sequences and cancer testes antigens have been suggested to be relatively late events in prostate cancer development [55,30]. In addition, metastatic sites have been reported to show greater divergence for DNA hypomethylation within some patients [30].

Studies showing clonal stability of DNA hypermethylation, and evidence of a subset of clones that are more closely related to metastatic disease, support the idea of using DNA methylation markers for prostate cancer detection and prognosis.

### 2.4. DNA methylation markers for the detection and stratification of prostate cancer

Combined panels of candidate DNA methylation markers have been shown to have high sensitivity and specificity for the discrimination of prostate cancers from benign tissue [51,38], with more recent studies showing proof of principle in prostate biopsy material [38]. Several studies have shown the potential for non-invasive monitoring of DNA methylation in cell-free DNA to detect prostate cancer [56–59]. Many of these studies show remarkable sensitivity and support the use of these tests for monitoring disease progression, however larger studies will be required to determine the clinical utility of these promising tests for prostate cancer diagnosis.

A molecular stratification for prostate cancer was proposed recently [28]. This large study found associations between genomic alterations and epigenetic profiles, that may represent a phenotypic difference between these molecular subtypes of prostate cancer. One third of ERG-positive tumours clustered together with a distinct hypermethylation profile [28]. The one percent of prostate cancers that harbour mutations in the gene encoding isocitrate dehydrogenase (*IDH1*) were found to have a divergent genome-wide hypermethylation profile [28]. This is likely to be a result of *IDH1*-R132H mutations driving production of the oncometabolite 2-hydroxyglutarate (2HG), with a reciprocal decrease in  $\alpha$ -ketoglutarate, a key component in the metabolic pathways of DNA methylation (the upstream metabolic processes are described in Section 3). It is currently unclear whether these molecular subtypes have clinical utility, what other factors impact on the observed



**Fig. 2.** Heatmap summary of genes and marker probes associated with DMRs reported in four or more studies. Left panel, correlations between gene expression (RNA-seq) and DNA methylation (450k arrays), from the PRAD TCGA data set (450k probe IDs indicated on the left). Middle panel, average methylation levels (Beta-values) from multiple studies spanning a range of prostate tissue types (450k probes indicated on the right). Right panel, androgen-stimulated gene expression changes in two prostate cancer cell lines (autocorrelation values denote a change with time following stimulation – a value of zero indicates no systematic change after stimulation). Missing data are indicated with a cross where not obvious.

methylation profiles, or what the phenotypic consequences of these epigenetic differences might be.

More effective diagnosis of prostate cancer would alleviate some of the current burden on health systems and decrease invasive procedures on healthy men. However, it is also critically important to distinguish indolent from aggressive prostate cancers so that aggressive treatments can be appropriately allocated to those patients who require such interventions, sparing other patients unnecessary over-treatment. Associations between DNA methylation changes and prognosis have been reported, including the correlation of *PTGS2* (*COX-2*), *HOXD3* and *ABHD9* hypermethylation with recurrence [51,22].

One study aiming to identify prognostic methylation markers for prostate cancer highlighted over one hundred candidate genomic loci [49]. However, the discrimination between relapsed and non-relapsed samples was far weaker than between tumour and benign samples [49], indicating more subtle differences between these groups. Among the candidate prognostic markers only *PTGS2* (*COX-2*) was validated from the previously mentioned studies. However, other studies have reported more promising findings by combining pilot genome-wide screening with targeted approaches on large test and validation cohorts [60]. A three gene methylation signature (*AOX1*, *C10RF114*, *HAPLN3*) was able to predict biochemical recurrence with respective hazard ratios of 1.9 and 2.3 in test and independent validation cohorts [60].

By taking a different approach and examining morphologically normal tissue adjacent to tumours, the methylation status of *GSTP1* and *APC* has been reported to have prognostic utility [61]. *GSTP1* and *APC* methylation in tumour adjacent tissue correlated with the methylation of these loci in matched tumour samples [61], consistent with either shared clonal ancestry or convergent evolution. Either hypothesis to explain these DNA methylation changes in tumour adjacent tissue would support a field-effect in a subset of

prostate cancers that may impact on outcome. These results are consistent with other reports of an epigenetic field effect (summarized in Section 2.2) and confirm other reports that epigenetic profiling could be a useful tool to avoid false-negatives in diagnostic biopsies [62].

In addition to these important targeted studies in large cohorts there is a need for methylation profiling studies that implement sensitive, genome-wide methods across samples representing the full range of prostate cancer disease stages to provide a clearer picture of the likely diagnostic and prognostic utility of these and other DNA methylation markers for prostate cancer.

### 2.5. Epigenetic regulation of alternative promoter usage in prostate cancer

In addition to gene silencing, DNA methylation can modulate gene isoform expression by impacting on alternative promoter regions. For example DNA methylation at the *RASSF1*, *APC* and *NDRG2* loci were shown to result in differential isoform expression [46]. This isoform selective expression was actively enforced by the epigenetic status at these loci, as shown by reversal of the isoform expression profile upon treatment with the demethylating agent 5-aza-cytidine [46]. Similar events have been reported in other studies in prostate cancer [63], however the true extent of this feature will only be made clear by large cohort studies combining comprehensive methylation profiling with RNA-sequencing on the same samples. An exemplar study leveraged the large RNA-sequencing gene expression data sets generated by TCGA groups to identify isoform switching in multiple tumour types (including prostate cancer), and speculate that epigenetic factors could be responsible [64]. In this study, tumour samples could be accurately identified solely by isoform switching signatures, highlighting the potential for isoform switching as a marker for prostate cancer.

## 2.6. Associations between epigenetic and genetic alterations in prostate cancer

Given the early and recurrent acquisition of *GSTP1* hypermethylation in prostate tumourigenesis many studies have suggested a role for *GSTP1* silencing in driving disease evolution by increasing the mutation rate [65–68]. Recent studies integrating methylation profiling with genome sequencing have uncovered additional associations between the epigenetic and genetic changes in prostate cancer.

Firstly, it has been shown that the methylation levels in matched benign samples are increased at mutated CpG sites in the tumour in comparison to non-mutated CpGs. [20]. This is consistent with the hypothesis that methylated cytosines are deaminated to uracil (and subsequently copied as thymine), a process believed to drive the observed high C-to-T mutation rates observed in prostate and other cancer types [69,5].

More surprisingly, it has been reported that sites of tandem duplication events in prostate cancer are frequently hypomethylated, while interchromosomal translocation break points are frequently hypermethylated [20]. These intriguing observations will need to be investigated in larger cohorts with paired methylome sequencing and genome sequencing to better characterize these associations.

Several studies have also suggested a link between *ETS* gene fusion status and DNA methylation profiles [46,45,48,28]. LINE repeats show differential methylation between ERG-positive and ERG-negative prostate tumours [46], differentially methylated regions associated with ERG status have been identified [48] and alternative mechanisms for *EZH2* activation have been proposed in ERG-negative tumours to phenocopy at least some of the consequences of *ETS* gene fusions [45]. The implications of molecular subtype differences in DNA methylation profiles are significant. Differences between *ETS*-fusion positive and negative cases highlight the interplay between epigenetic state, gene rearrangements and hormone signalling, since the AR regulates *ETS*-fusions and AR signalling is altered in *ETS*-positive tumours [15]. Equally, these effects could lead to misinterpretation of studies that cannot take them into account.

## 2.7. Epigenetic changes impacting on hormone signalling in prostate cancer

In addition to interactions with genomic events, the epigenetic profile has been linked to AR signalling, both as a modulator of hormone response and a driver event in progression to Castration Resistant Prostate Cancer (CRPC). AR-bound enhancers were observed to show greater intratumoural DNA methylation variation than other enhancer sites [27], suggesting clonal plasticity in the AR regulome.

An integrative analysis of copy number and DNA methylation in CRPC revealed convergence on the androgen synthesis pathway, with copy number and methylation changes converging on *HSD17B2* and other targets that may be involved in hormone therapy resistance [10].

Loss of AR protein expression is a characteristic of hormone-relapsed prostate cancer (PCa) that is no longer dependant on AR signalling, a phenotype that is becoming more common following the clinical use of second generation AR targeting therapies [70,71]. Epigenetic silencing of *AR* gene expression has been reported in prostate tumour samples [72,73] and treatment of AR negative PCa cells with the global demethylating agent 5-aza-deoxycytidine can induce functional AR signalling in these cells [73–76]. However, other studies in CRPC have reported no change in AR promoter methylation [10], suggesting either differences between analysis

methods or possibly patient cohorts treated with different hormone therapies.

Combining androgen signalling blockade with the demethylating agent 5-aza-deoxycytidine increased response in the preclinical models of PCa [77,75], suggesting that future studies combining demethylating agents with second generation AR blockade may improve patient outcome or delay relapse by targeting the AR, its target genes or upstream pathways.

Oestrogen receptor alpha and beta methylation have also been identified in some studies of prostate cancer samples [78,51,46,26]. Although these methylation changes at *ER* genes are not highlighted in the majority of studies it is noteworthy that studies reporting *ER* gene methylation used MSP [51], MBD-pulldown [26] or restriction enzyme based methods [46] (consistent with either limitations of the more commonly used 450k arrays at these loci or convergent artefacts between the other methods). Future studies should aim to assay DNA methylation at the *ERS1* and *ESR2* loci in prostate tissues to determine whether this could affect the interplay between nuclear hormone receptors and be another mechanism through which epigenetic changes impact on hormone signalling in prostate cancer.

A large number of these putative epigenetic markers were identified by methylation profiling of prostate cancer cell lines. In addition these cell line models are the most widely used tools for functional validation studies of candidate gene silencing or derepression events identified from genome-wide profiling studies. Therefore we summarize the utility and limitations of these models below.

## 2.8. Prostate cancer cell lines


It has for some time been recognized that immortalized cell lines do not, in respect of their methylation patterns, reflect the cells from which they originate; typically immortalized cells exhibit hypermethylation of CpG islands [79–81] although it has been noted that this may represent selection pressure where highly methylated cells are more likely to be immortalized successfully [82] and there are suggestions that the methylation changes can predate immortalization [83].

The characteristic methylation changes that occur in immortalized cells are similar to the changes seen in cancers [84,85] raising the hope that the epigenetics of cancer cell lines may represent malignant tissue well. Recent reports argue that it is specifically the immortalization of cells rather than other oncogenic activity that leads to changes in methylation profile [86]. Intriguingly, sites that are methylated in cancer cell lines are enriched for NANOG binding sites [87]. As well as its role in maintaining stem cell pluripotency, NANOG has been shown to be pro-tumourigenic in prostate cancer cell lines, conferring cancer-stem-cell-like properties [88], and is directly androgen regulated [89].

Early targeted studies identified good agreement between primary cancers and cell lines [90,91], but broader differences may mean that only a minority of tumours are well-represented by cell lines. Genome-wide profiling has revealed xenografts to be a better model in head-and-neck squamous cell carcinomas [92], while a recent paper has reported near-complete loss of 5-hydroxymethylcytosine [93] – raising questions about cell lines' value in this regard.

The utility of prostate cell lines is further affected by the fact that the commonly used prostate cancer cell line models (recently reviewed elsewhere [94]) were derived from metastases, and metastasis itself is associated with changes in methylation profiles [95,96]. Nevertheless, substantial public data exist for a number of prostate cell lines as denoted in Table 1, notably LNCaP, PC3 DU-145 and PrEC, making them an attractive resource.





**Fig. 3.** Cell lines as a representation of primary tumours. (A) Clustering of our example cell line and tissue data sets (described in Section 1.3). The distance between two samples is defined as Cohen's Kappa measure of agreement (applied after dichotomizing methylation beta values). (B) Illustrating, from the cell line and tissue data sets, the median proportion of methylation at loci near the *GSTP1* promoter. As expected, the cancer cell lines are generally hypermethylated in this region, as are the neoplastic and malignant tissues. (C) Illustrating, from the cell line and tissue data sets, the median proportion of methylation at loci near the *TERT* gene. An area in the gene promoter shows progressively increased methylation levels, with the cancer cell lines' levels most in keeping with metastases.

In Fig. 3A, we cluster the available Illumina Infinium Human Methylation450 BeadChip cell line data with our example Tissue data set. It is notable that inter-sample heterogeneity increases as one progresses through normal, benign, tumour and metastasis samples. Reflecting their origins, the prostate cancer cell lines are more alike the metastasis samples than they are the primary tumour samples, while the PrEC cells cluster with the normal samples.

It should be noted that all of the cell lines show levels of agreement with primary tumours that are substantially above chance, and that their greater similarity to metastases only requires care to be taken over the interpretation of any results arising. The cell lines still reflect the behaviour of primary tumours at key loci such as *GSTP1* (Fig. 3B), although at loci such as the promoter of *TERT*

there appears to be progressively greater DNA methylation as one moves from normal tissue, through hyperplastic and neoplastic tissues to primary tumours and metastases, the malignant cell lines showing greater values still (Fig. 3C).

Despite the inevitable caveats about the use of cell lines, they offer natural advantages for the inference of function. They enable one to run controlled experiments with identical subjects in each/every arm, and make it possible to measure multiple characteristics (e.g. genome-wide methylation and transcript abundance) on effectively the same samples. Both of these have been exploited to address questions of prostate cancer biology.

To understand better the methylation-driven regulation of the cancer genome, mRNA expression data are the natural orthogonal data to bring into an integrative analysis. In this manner, the methylation-regulated nature of key genes and alternative transcriptional start site usage have been explored in prostate cancer [46,97].


A substantial body of work has linked DNA methylation with other epigenetic marks better to understand gene regulation in (prostate) cancer. The H3K27me3 mark at promoters is associated with inactivated genes, while methylation of promoters is similarly associated with gene repression, but the two mechanisms have been seen to be neither exclusive nor deterministically linked in LNCaP [98]. Where the two mechanisms do coexist, a dual therapy to reinstate expression of tumour suppressor genes becomes a possibility [99]. Elsewhere it has been shown that a genome-wide restructuring of nucleosome densities is associated with changes in DNA methylation of enhancer regions of PCa cell lines [100].

Other studies have sought to explain epigenetic mechanisms of treatment and resistance to treatment. Epigenetic silencing of *SLFN11* has been associated with resistance to platinum-based chemotherapies in a number of cell lines including DU-145 and PC3 [101], an epigenetic mechanism of the preventative agents sulforaphane and 3,3'-diindolylmethane is elucidated [102], and the mechanism of Genistein has been shown not to be dependent on broad methylation changes, but rather histone acetylation [103].

Perhaps most interestingly for our topic, a recent report has shown that dosing cells with S-adenosylmethionine (SAM) inhibits invasion [104]; the same group having previously identified a role for hypomethylation in the metastasis of prostate cancer [105]. SAM is an important methyl donor for histone, DNA and RNA methylation, and all general protein lysine and arginine methylation (as described in Section 3). The demonstration, in vitro and in vivo, that replenishing the reservoir of methyl donors can inhibit metastasis (while only increasing the methylation status of specific loci rather than reversing the genome-wide hypomethylation) provides some evidence of a mechanistic role for the methylation patterns in metastases, highlights the importance of the available metabolic pool for cancer progression, and invites consideration of the broader role of SAM in the metabolic pathways.

### 3. Effects of the metabolic pool on DNA methylation

The prostate gland is a metabolically specialized organ responsible for supporting sperm viability. This specialization is characterized by net secretion of citrate into the seminal fluid [106]. Whilst a significant reduction in the production of these metabolites has been reported in numerous magnetic resonance spectroscopy studies on clinical samples, the molecular drivers for this down-regulation remain controversial [107–110]. Of the polyamines normally produced by the prostate gland, spermine is particularly abundant [111]. Rat models of castration-induced regression and testosterone-induced regrowth of the prostate gland have shown that polyamine production is




**Fig. 4.** Schematic representation of metabolic pathways that impact on DNA methylation in prostate cancer. The core DNA methylation pathway is represented at the top of the schematic, with connected pathways of relevance to prostate cancer shown below. Colour key relates to metadata annotations shown in Fig. 5. Values for each metabolic step are compressed into a single summary to simplify the schematic. Metabolic steps are annotated if: androgen regulation in LNCaP or VCaP gives autocorrelation values >0.5; there is a mean methylation change between tumour and normal tissue of  $\pm 0.2$ ; the correlation of methylation and expression changes is >0.3 or <-0.3 (see Fig. 5 for details).

tightly regulated by androgens, in part through control over the expression of key biosynthetic enzymes (ornithinedecarboxylase (ODC), S-adenosylmethionine decarboxylase (AMD1) and spermidine synthase (SMS); highlighted in Figs. 4 and 5) [112–114]. The methyl donor in this pathway is S-adenosylmethionine (decarboxy-AdoMet), a metabolite that is directly downstream of the DNA methylation donor S-adenosylmethionine (SAM, AdoMet). Therefore, alterations in the flux through either polyamine metabolism or DNA methylation would be expected to affect the available pools of methyl-donors, with reciprocal effects on the flux of the other pathway.

The prostate has one of the highest concentrations of polyamines of any tissue in the body and the expression of these enzymes is associated with glandular epithelial cells with significant quantities of polyamines secreted into seminal fluids. Spermine levels in clinical samples have been reported to correlate positively with the differentiation status of the tumour and in preclinical models to promote growth inhibition [115,116].

Polyamine synthesis requires one-carbon metabolism and in particular methionine metabolism with S-adenosylmethionine (SAM) which upon decarboxylation acts as the primary amino-propyl donor for polyamine synthesis (Fig. 4) [117]. Consequently the fate of methionine and its derivatives may be influenced by



**Fig. 5.** Heatmap summary of genes in metabolic pathways that impact on DNA methylation (relating to Fig. 4). Left panel, correlations between gene expression (RNA-seq) and DNA methylation (450k arrays), from the PRAD TCGA data set (450k probe IDs indicated on the left). Middle panel, average methylation levels (Beta-values) from multiple studies spanning a range of prostate tissue types (450k probes indicated on the right). Right panel, androgen-stimulated gene expression changes in two prostate cancer cell lines (autocorrelation values denote a change with time following stimulation).

changes in the specialized secretory functions of the prostate gland as cancer emerges, in particular a decline in polyamine biosynthesis and secretion may enhance the pool of SAM available to support epigenetic modifications (metabolic pathway links shown in Fig. 4). Proving causative associations between rates of polyamine biosynthesis and regulation of the epigenetic state via the availability of SAM is extremely challenging, since methylation patterns can be highly locus and cell-type dependent and are supported by a complex regulatory network downstream of metabolite consumption and upstream of DNA substrates. Furthermore, one-carbon metabolism consists of a number of additional interconnected metabolic processes that may impinge on polyamines and the epigenome (folate cycle, methionine cycle and glycine/serine metabolism, Fig. 4).

Prostate cancer is characterized by the activity of transcription factors, particularly AR but also others such as c-Myc. A natural question then is how the transcription factors and important enzymes interact with the metabolome and epigenetic status of a tumour. As indicated earlier, the synthesis of polyamines and the expression of the key enzymes required for this and a number of other metabolic processes upstream of SAM production are driven by AR and associated with differentiated prostate cancers (Figs. 4 and 5). Glycine N-methyltransferase (GNMT) is an enzyme which converts glycine to sarcosine and in the process converts SAM to S-adenosylhomocysteine (Fig. 4). In cell-lines expression of

the GNMT enzyme is androgen dependent and in tissue samples it has been shown to be over-expressed in prostate cancers (Figs. 4 and 5) [118,40]. Sarcosine, the product of the reaction catalysed by GNMT, has also been reported to be elevated and detectable in urine samples and some studies have associated this with the emergence of castrate-resistant disease [119,120]. Of the enzymes involved in one-carbon metabolism GNMT is therefore currently the most extensively characterized androgen-dependent component of the pathway. By contrast the expression of enzymes required for serine metabolism appears in prostate cancer cell-lines to be repressed by androgens and in other cancer models to be induced by c-Myc [40,121]. Whilst serine metabolism also impacts on the methionine cycle, the most direct impact of serine consumption appears to be to sustain de novo nucleotide biosynthesis in support of elevated rates of DNA replication and/or transcription and cell proliferation [122]. This in turn is often a feature of cell cycle dysregulation, characteristic of late-stage, metastatic prostate cancer [123]. Hypothetically a phasic transition in one-carbon metabolism may therefore consist initially of reduced rates of polyamine biosynthesis and enhanced DNA and histone methylation in localized disease transitioning into enhanced serine metabolism during metastatic progression. This would at least be consistent with the observations that DNA hypermethylation is an earlier event in prostate tumourigenesis, while hypomethylation may occur in later stage disease (as described in Section 2).

The DNA methylation status of a subset of genes encoding these metabolic enzymes increases between normal samples compared to benign, tumour and metastasis, consistent with early hypermethylation changes (Fig. 5). A separate set of enzyme-encoding genes show hypomethylation in metastatic samples compared to localized and pre-malignant samples (Fig. 5), again consistent with genome-wide observations of later stage hypomethylation (Section 2). It is of interest that there appears to be exclusivity of regulation within the metabolism-related genes illustrated here (Fig. 5), with AR-regulated genes showing little evidence of differential methylation, and the strongest differentially methylated genes not being AR-regulated. This despite the independence of the data sets from which these characteristics were identified.

In addition to the potential influence of upstream enzyme expression and metabolite pools on DNA methylation, the demethylation pathway is intricately linked to central metabolism. The TCA cycle metabolite alpha-ketoglutarate is required for TET enzyme activity, the first step in cytosine demethylation (Fig. 4). Therefore, mitochondrial function may have a profound effect on both cytosine methylation and the levels of hydroxymethylation in the genome of a cell.

#### 4. Mitochondrial methylation and prostate cancer

##### 4.1. Importance of mitochondria in PCA

While of general interest in cancer (reviewed, for example by Wallace [124]), mitochondria are of specific interest in prostate cancer beyond the characteristic zinc-inhibition of TCA cycle and AR regulation of certain metabolites described in the previous section. Recent 'pan-cancer' analyses have identified many mutations in mitochondrial DNA, without showing them to be drivers [125,126], but animal models have demonstrated the functional impact of mitochondrial genetic mutations on prostate cancer [127] and a cohort study in humans has identified associations with proliferation [3].

The case for DNA methylation changes in the prostate cancer mitochondria themselves is not as clear cut as that for genetic changes. Methylation of mitochondrial DNA has been controversial since it was first reported [128]. Subsequent papers variously confirmed this result [129] or reported no such methylation [130]. The discussion has continued since with arguments for and against based on sequence analysis [131,132], and use of methylation sensitive and insensitive restriction enzymes [133,134]. One aspect of the argument against the existence of methylation in the mitochondria has been the absence of the actors that facilitate methylation of nuclear DNA. However, recent years have seen the identification of


methyl donors [135], methyltransferases [136,137], and even TET1 present in the mitochondria [138]. This latter observation supporting reports of 5-hydroxymethylcytosine in the mitochondrial DNA [136,139]. The history of this topic is reviewed more thoroughly elsewhere [140].

Recently there have been studies that provide stronger evidence that epigenetic modifications of mitochondrial DNA do indeed take place. For example, one study identified methylated bases in nucleus-free platelets [141], while another used orthogonal and complementary technologies to profile the mitochondrial methylation in a wide range of tissue and cell types [142]. Given the unique behaviour of mitochondria in prostate cancer, if mitochondrial DNA can be methylated, this is an aspect that may reward investigation.

##### 4.2. Reciprocal regulation of nuclear methylation and mitochondria

Apart from direct epigenetic changes to the mtDNA, the mitochondria have a complex relationship with epigenetic alterations to the nuclear DNA. Of the order of a thousand coding genes have products that are active in the mitochondria, and only 13 originate from the mitochondrial DNA. It follows that any epigenetic regulation of the remaining mitochondrial actors in the nucleus will likely influence mitochondrial behaviour. One study identified tissue-specific differentially methylated regions in mitochondrial-acting nuclear-encoded genes [143] while another has concluded that epigenetic regulation of mitochondrial-acting nuclear-encoded genes was higher than other nuclear-encoded genes [144]. This latter result can be replicated in prostate cancer using a single tumour sample (Fig. 6A). While this is representative of the other samples in our example set, with >5000 probes from regions around the transcription start sites of mitochondrial actors showing consistent hypomethylation and approximately 400 showing consistent hypermethylation, there is also evidence of changes in the regulation of these regions with the progression of disease. Fig. 6B shows a heatmap of the 114 most variable probes in these regions, and it is apparent that again there are clusters of probes that variously gain methylation in neoplastic tissue and continue to do so in tumour and metastases, others with the opposite pattern, and smaller numbers where the changes are primarily defined by the metastases samples.

DNA polymerase gamma is responsible for the replication of mitochondrial DNA and is regulated by the methylation of the *POLG* gene [145,146] leading to associations between mtDNA copy number and *POLG* methylation levels. Methylation of the *PPARGC1A* gene has also been seen to correlate negatively with mtDNA copy number.



**Fig. 6.** (A) Reprod. Fig. 4 of Chinnery et al. for a high-cellularity prostate cancer sample. Probes on the Infinium HumanMethylation450 BeadChip that lay within 1000 bases of transcription start sites (TSS) were divided between genes listed in MitoCarta and those that are not. (B) A heatmap of median beta methylation values for different tissues across probes that show most variation in our data.



While the methylation of some genes can regulate mtDNA copy number, it has also been shown that mtDNA copy number can regulate some nuclear genes [144,147]. Furthermore, studies using cybrids have shown that the mitochondria can affect nuclear methylation patterns [139,148]. Given the importance of epigenetics to prostate cancer regulation, and the characteristic behaviour of mitochondria (and broader metabolic pathways) in the disease, these associations demand attention in attempts to unravel prostate cancer biology.

## 5. Current perspectives and future directions for the role of epigenomic changes in prostate cancer

It is clear that clonal expansions of cells with stable epigenomic changes occur in prostate cancer. DNA methylation changes are the most recurrent events so far identified in prostate cancer, and specific changes may associate with outcome. The epigenome continues to evolve throughout the life history of prostate cancer, with distinct features presenting at different stages and interacting with specific genomic changes. It will be crucial to overlay other epigenetic changes within the same cohorts of samples to build up a picture of the epigenetic landscape in prostate cancer. Future studies are also needed to integrate both genomic and epigenomic data in large cohorts of samples, to elucidate the interaction between genomic and epigenetic changes, to provide a more comprehensive view of the pathways affected in each prostate tumour sample, and to assess clinical associations with specific sets of changes.

## Acknowledgements

CEM is funded by an ERC grant. IGM is supported in Oslo by funding from the Norwegian Research Council, Helse Sor-Ost and the University of Oslo through the Centre for Molecular Medicine (Norway), which is a part of the Nordic EMBL (European Molecular Biology Laboratory) partnership. IGM holds a visiting scientist position with Cancer Research UK through the Cambridge Research Institute and a Senior Honorary Visiting Research Fellowship with Cambridge University through the Department of Oncology. IGM is supported in Belfast by the Belfast-Manchester Movember Centre of Excellence (CE013.2-004), funded in partnership with Prostate Cancer UK. AGL is supported by a Cancer Research UK programme grant (C14303/A20406) to Simon Tavaré and by the European Commission through the Horizon 2020 project SOUND (Grant Agreement no. 633974). CEM and AGL acknowledge the support of the University of Cambridge, Cancer Research UK and Hutchison Whampoa Limited.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jsbmb.2016.04.009>.

## References

- [1] C.E. Barbieri, S.C. Baca, M.S. Lawrence, F. Demichelis, M. Blattner, J.-P. Theurillat, T.A. White, P. Stojanov, E. Van Allen, N. Stransky, et al., Exome sequencing identifies recurrent *spop*, *foxa1* and *med12* mutations in prostate cancer, *Nat. Genet.* 44 (2012) 685–689.
- [2] M. Kobayashi, H. Ishida, T. Shindo, S.-I. Niwa, M. Kino, K. Kawamura, N. Kamiya, T. Imamoto, H. Suzuki, Y. Hirokawa, T. Shiraishi, T. Tanizawa, Y. Nakatani, T. Ichikawa, Molecular analysis of multi-focal prostate cancer by comparative genomic hybridization, *Prostate* 68 (2008) 1715–1724.
- [3] J. Lindberg, I.G. Mills, D. Klevebring, W. Liu, M. Neiman, J. Xu, P. Wikström, P. Wiklund, F. Wiklund, L. Egevad, H. Grönberg, The mitochondrial and autosomal mutation landscapes of prostate cancer, *Eur. Urol.* 63 (2013) 702–708.
- [4] L.K. Boyd, X. Mao, L. Xue, D. Lin, T. Chaplin, S.C. Kudahetti, E. Stankiewicz, Y. Yu, L. Beltran, G. Shaw, J. Hines, R.T.D. Oliver, D.M. Berney, B.D. Young, Y.-J. Lu, High-resolution genome-wide copy-number analysis suggests a monoclonal origin of multifocal prostate cancer, *Genes Chromosomes Cancer* 51 (2012) 579–589.
- [5] C.S. Cooper, R. Eeles, D.C. Wedge, P. Van Loo, G. Gundem, L.B. Alexandrov, B. Kremeyer, A. Butler, A.G. Lynch, N. Camacho, et al., Analysis of the genetic phylogeny of multifocal prostate cancer identifies multiple independent clonal expansions in neoplastic and morphologically normal prostate tissue, *Nat. Genet.* 47 (2015) 367–372.
- [6] G. Gundem, P. Van Loo, B. Kremeyer, L.B. Alexandrov, J.M. Tubio, E. Papaemmanuil, D.S. Brewer, H.M. Kallio, G. Högn&rdquo;as, M. Annala, et al., The evolutionary history of lethal metastatic prostate cancer, *Nature* (2015).
- [7] H. Ross-Adams, A. Lamb, M. Dunning, S. Halim, J. Lindberg, C. Massie, L. Egevad, R. Russell, A. Ramos-Montoya, S. Vowler, et al., Integration of copy number and transcriptomics provides risk stratification in prostate cancer: a discovery and validation cohort study, *EBioMedicine* 2 (2015) 1133–1144.
- [8] S.A. Tomlins, D.R. Rhodes, S. Perner, S.M. Dhanasekaran, R. Mehra, X.-W. Sun, S. Varambally, X. Cao, J. Tchinda, R. Kuefer, et al., Recurrent fusion of *TMPS2* and *ETS* transcription factor genes in prostate cancer, *Science* 310 (2005) 644–648.
- [9] W.-H. Lee, W.B. Isaacs, G.S. Bova, W.G. Nelson, CG island methylation changes near the *GSTP1* gene in prostatic carcinoma cells detected using the polymerase chain reaction: a new prostate cancer biomarker, *Cancer Epidemiol. Biomarkers Prev.* 6 (1997) 443–450.
- [10] T.W. Friedlander, R. Roy, S.A. Tomlins, V.T. Ngo, Y. Kobayashi, A. Azameera, M.A. Rubin, K.J. Pienta, A. Chinnaiyan, M.M. Ittmann, et al., Common structural and epigenetic changes in the genome of castration-resistant prostate cancer, *Cancer Res.* 72 (2012) 616–625.
- [11] M.T. Bedford, P.D. Van Helden, Hypomethylation of DNA in pathological conditions of the human prostate, *Cancer Res.* 47 (1987) 5274–5276.
- [12] W.-H. Lee, R.A. Morton, J.I. Epstein, J.D. Brooks, P.A. Campbell, G.S. Bova, W.-S. Hsieh, W.B. Isaacs, W.G. Nelson, Cytidine methylation of regulatory sequences near the *pi*-class glutathione S-transferase gene accompanies human prostatic carcinogenesis, *Proc. Natl. Acad. Sci. U.S.A.* 91 (1994) 11733–11737.
- [13] J.G. Herman, A. Merlo, L. Mao, R.G. Lapidus, J.-P.J. Issa, N.E. Davidson, D. Sidransky, S.B. Baylin, Inactivation of the *CDKN2/p16/MTS1* gene is frequently associated with aberrant DNA methylation in all common human cancers, *Cancer Res.* 55 (1995) 4525–4530.
- [14] N. Ahmadiyeh, M.M. Pomerantz, C. Grisanzio, P. Herman, L. Jia, V. Almendro, H.H. He, M. Brown, X.S. Liu, M. Davis, et al., 8q24 prostate, breast, and colon cancer risk loci show tissue-specific long range interaction with *MYC*, *Proc. Natl. Acad. Sci. U.S.A.* 107 (2010) 9742–9746.
- [15] J. Yu, J. Yu, R.-S. Mani, Q. Cao, C.J. Brenner, X. Cao, X. Wang, L. Wu, J. Li, M. Hu, et al., An integrated network of androgen receptor, polycomb, and *TMPS2-ERG* gene fusions in prostate cancer progression, *Cancer Cell* 17 (2010) 443–454.
- [16] K. Xu, Z.J. Wu, A.C. Groner, H.H. He, C. Cai, R.T. Lis, X. Wu, E.C. Stack, M. Loda, T. Liu, et al., *EZH2* oncogenic activity in castration resistant prostate cancer cells is polycomb-independent, *Science* 338 (2012) 1465–1469.
- [17] H. Yang, Y. Liu, F. Bai, J. Zhang, S. Ma, J. Liu, Z. Xu, H. Zhu, Z. Ling, D. Ye, et al., Tumor development is associated with decrease of *TET* gene expression and 5-methylcytosine hydroxylation, *Oncogene* 32 (2013) 663–669.
- [18] R. Lister, M. Pelizzola, R.H. Dowen, R.D. Hawkins, G. Hon, J. Tonti-Filippini, J.R. Nery, L. Lee, Z. Ye, Q.-M. Ngo, et al., Human DNA methylomes at base resolution show widespread epigenomic differences, *Nature* 462 (2009) 315–322.
- [19] F.E. Garrett-Bakelman, C.K. Sheridan, T.J. Kacmarczyk, J. Ishii, D. Betel, A. Alonso, C.E. Mason, M.E. Figueroa, A.M. Melnick, Enhanced reduced representation bisulfite sequencing for assessment of DNA methylation at base pair resolution, *J. Vis. Exp.* 96 (2015) e52246.
- [20] P.-C. Lin, E.G. Giannopoulou, K. Park, J.M. Mosquera, A. Sboner, A.K. Tewari, L.A. Garraway, H. Beltran, M.A. Rubin, O. Elemento, Epigenomic alterations in localized and advanced prostate cancer, *Neoplasia* 15 (2013) 373–383.
- [21] J. Sandoval, H. Heyn, S. Moran, J. Serra-Musach, M.A. Pujana, M. Bibikova, M. Esteller, Validation of a DNA methylation microarray for 450,000 CpG sites in the human genome, *Epigenetics* 6 (2011) 692–702.
- [22] M. Stott-Miller, S. Zhao, J.L. Wright, S. Kolb, M. Bibikova, B. Klotzle, E.A. Ostrander, J.-B. Fan, Z. Feng, J.L. Stanford, Validation study of genes with hypermethylated promoter regions associated with prostate cancer recurrence, *Cancer Epidemiol. Biomarkers Prev.* 23 (2014) 1331–1339.
- [23] M. Weber, J.J. Davies, D. Wittig, E.J. Oakeley, M. Haase, W.L. Lam, D. Schuebeler, Chromosome-wide and promoter-specific analyses identify sites of differential DNA methylation in normal and transformed human cells, *Nat. Genet.* 37 (2005) 853–862.
- [24] J.-H. Luo, Y. Ding, R. Chen, G. Michalopoulos, J. Nelson, G. Tseng, P.Y. Yan, Genome-wide methylation analysis of prostate tissues reveals global methylation patterns of prostate cancer, *Am. J. Pathol.* 182 (2013) 2028–2036.
- [25] E.U. Selker, N.A. Tountas, S.H. Cross, B.S. Margolin, J.G. Murphy, A.P. Bird, M. Freitag, The methylated component of the *Neurospora crassa* genome, *Nature* 422 (2003) 893–897.
- [26] M.J. Aryee, W. Liu, J.C. Engelmann, P. Nuhn, M. Gurel, M.C. Håner, D. Esopi, R.A. Irizarry, R.H. Getzenberg, W.G. Nelson, J. Luo, J. Xu, W.B. Isaacs, G.S. Bova, S. Yegnasubramanian, DNA methylation alterations exhibit intraindividual stability and interindividual heterogeneity in prostate cancer metastases, *Sci. Transl. Med.* 5 (2013) 169ra10.
- [27] D. Brooks, Y. Assenov, S. Minner, O. Bogatyrova, R. Simon, C. Koop, C. Oakes, M. Zucknick, D.B. Lipka, J. Weischenfeldt, et al., Intratumor DNA methylation



- heterogeneity reflects clonal evolution in aggressive prostate cancer, *Cell Rep.* 8 (2014) 798–806.
- [28] Cancer Genome Atlas Research Network, The molecular taxonomy of primary prostate cancer, *Cell* 163 (2015) 1011–1025.
- [29] A.C. Rand, M. Jain, J. Eizenga, A. Musselman-Brown, H.E. Olsen, M. Akeson, P. Benedict, Cytosine variant calling with high-throughput nanopore sequencing, *bioRxiv* (2016), <http://dx.doi.org/10.1101/047134>.
- [30] S. Yegnasubramanian, M.C. Häner, Y. Zhang, B. Gurel, T.C. Cornish, Z. Wu, R.A. Irizarry, J. Morgan, J. Hicks, T.L. DeWeese, et al., DNA hypomethylation arises later in prostate cancer progression than CpG island hypermethylation and contributes to metastatic tumor heterogeneity, *Cancer Res.* 68 (2008) 8954–8967.
- [31] R.A. Irizarry, C. Ladd-Acosta, B. Carvalho, H. Wu, S.A. Brandenburg, J.A. Jeddloh, B. Wen, A.P. Feinberg, Comprehensive high throughput arrays for relative methylation (CHARM), *Genome Res.* 18 (2008) 780–790.
- [32] G. Landan, N.M. Cohen, Z. Mukamel, A. Bar, A. Molchadsky, R. Brosh, S. Horn-Saban, D.A. Zalcenstein, N. Goldfinger, A. Zundevich, et al., Epigenetic polymorphism and the stochastic formation of differentially methylated regions in normal and cancerous tissues, *Nat. Genet.* 44 (2012) 1207–1214.
- [33] D.A. Landau, K. Clement, M.J. Ziller, P. Boyle, J. Fan, H. Gu, K. Stevenson, C. Sougnez, L. Wang, S. Li, et al., Locally disordered methylation forms the basis of intratumor methylome variation in chronic lymphocytic leukemia, *Cancer Cell* 26 (2014) 813–825.
- [34] K.D. Siegmund, P. Marjoram, Y.-J. Woo, S. Tavaré, D. Shibata, Inferring clonal expansion and cancer stem cell dynamics from DNA methylation patterns in colorectal cancers, *Proc. Natl. Acad. Sci. U.S.A.* 106 (2009) 4828–4833.
- [35] K.D. Hansen, W. Timp, H.C. Bravo, S. Sabuncian, B. Langmead, O.G. McDonald, B. Wen, H. Wu, Y. Liu, D. Diep, et al., Increased methylation variation in epigenetic domains across cancer types, *Nat. Genet.* 43 (2011) 768–775.
- [36] A. Diez-Villanueva, I. Mallona, M.A. Peinado, Wanderer, an interactive viewer to explore DNA methylation and gene expression data in human cancer, *Epigenet. Chromatin* 8 (2015) 1.
- [37] H. Naem, N.C. Wong, Z. Chatterton, M.K.H. Hong, J.S. Pedersen, N.M. Corcoran, C.M. Hovens, G. Macintyre, Reducing the risk of false discovery enabling identification of biologically significant genomewide methylation status using the HumanMethylation450 array, *BMC Genomics* 15 (2014) 51.
- [38] A. Paziewska, M. Dabrowska, K. Goryca, A. Antoniewicz, J. Dobruch, M. Mikula, D. Jarosz, L. Zapala, A. Borowka, J. Ostrowski, DNA methylation status is more reliable than gene expression at detecting cancer in prostate biopsy, *Br. J. Cancer* 111 (2014) 781–789.
- [39] C.E. Massie, I. Spiteri, H. Ross-Adams, H. Luxton, J. Kay, H.C. Whitaker, M.J. Dunning, A.D. Lamb, A. Ramos-Montoya, D.S. Brewer, et al., HES5 silencing is an early and recurrent change in prostate tumorigenesis, *Endocr.-Relat. Cancer* 22 (2015) 131–144.
- [40] C.E. Massie, A. Lynch, A. Ramos-Montoya, J. Boren, R. Stark, L. Fazli, A. Warren, H. Scott, B. Madhu, N. Sharma, et al., The androgen receptor fuels prostate cancer by regulating central metabolism and biosynthesis, *EMBO J.* 30 (2011) 2719–2733.
- [41] D.S. Millar, K.K. Ow, C.L. Paul, P.J. Russell, P.L. Molloy, S.J. Clark, et al., Detailed methylation analysis of the glutathione S-transferase  $\pi$  (GSTP1) gene in prostate cancer, *Oncogene* 18 (1999) 1313–1324.
- [42] S. Santouridis, A. Flori, R. Ackermann, H.-C. Wirtz, W.A. Schulz, High frequency of alterations in DNA methylation in adenocarcinoma of the prostate, *Prostate* 39 (1999) 166–174.
- [43] P.J. Bastian, S. Yegnasubramanian, G.S. Palapattu, C.G. Rogers, X. Lin, A.M. De Marzo, W.G. Nelson, Molecular biomarker in prostate cancer: the role of CpG island hypermethylation, *Eur. Urol.* 46 (2004) 698–708.
- [44] R.E. Jimenez, A.H. Fischer, J.A. Petros, M.B. Amin, Glutathione S-transferase  $\pi$  gene methylation: the search for a molecular marker of prostatic adenocarcinoma, *Adv. Anat. Pathol.* 7 (2000) 382–389.
- [45] S.T. Börno, A. Fischer, M. Kerick, M. F&rdquo;alth, M. Laible, J.C. Brase, R. Kuner, A. Dahl, C. Grimm, B. Sayanjali, et al., Genome-wide DNA methylation events in TMPRSS2-ERG fusion-negative prostate cancers implicate an EZH2-dependent mechanism with miR-26a hypermethylation, *Cancer Discov.* 2 (2012) 1024–1035.
- [46] J.H. Kim, S.M. Dhanasekaran, J.R. Prensner, X. Cao, D. Robinson, S. Kalyana-Sundaram, C. Huang, S. Shankar, X. Jing, M. Iyer, et al., Deep sequencing reveals distinct patterns of DNA methylation in prostate cancer, *Genome Res.* 21 (2011) 1028–1041.
- [47] K. Kron, V. Pethe, L. Briollais, B. Sadikovic, H. Ozelik, A. Sunderji, V. Venkateswaran, J. Pinthus, N. Fleshner, T. van der Kwast, et al., Discovery of novel hypermethylated genes in prostate cancer using genomic CpG island microarrays, *PLoS ONE* 4 (2009) e4830.
- [48] K. Kron, D. Trudel, V. Pethe, L. Briollais, N. Fleshner, T. van der Kwast, B. Bapat, Altered DNA methylation landscapes of polycomb-repressed loci are associated with prostate cancer progression and ERG oncogene expression in prostate cancer, *Clin. Cancer Res.* 19 (2013) 3450–3461.
- [49] S. Mahapatra, E.W. Klee, C.Y. Young, Z. Sun, R.E. Jimenez, G.G. Klee, D.J. Tindall, K.V. Donkena, Global methylation profiling for risk prediction of prostate cancer, *Clin. Cancer Res.* 18 (2012) 2882–2895.
- [50] B. Yang, S. Bhusari, J. Kueck, P. Weerathunga, J. Wagner, G. Levenson, W. Huang, D.F. Jarrard, Methylation profiling defines an extensive field defect in histologically normal prostate tissues associated with prostate cancer, *Neoplasia* 15 (2013) 399–408.
- [51] S. Yegnasubramanian, J. Kowalski, M.L. Gonzalgo, M. Zahurak, S. Piantadosi, P.C. Walsh, G.S. Bova, A.M. De Marzo, W.B. Isaacs, W.G. Nelson, Hypermethylation of CpG islands in primary and metastatic human prostate cancer, *Cancer Res.* 64 (2004) 1975–1986.
- [52] Y.P. Yu, S. Paranjpe, J. Nelson, S. Finkelstein, B. Ren, D. Kokkinakis, G. Michalopoulos, J.-H. Luo, High throughput screening of methylation status of genes in prostate cancer using an oligonucleotide methylation array, *Carcinogenesis* 26 (2005) 471–479.
- [53] J.D. Brooks, M. Weinstein, X. Lin, Y. Sun, S.S. Pin, G.S. Bova, J.I. Epstein, W.B. Isaacs, W.G. Nelson, Cg island methylation changes near the *gstp1* gene in prostatic intraepithelial neoplasia, *Cancer Epidemiol. Biomarkers Prev.* 7 (1998) 531–536.
- [54] A. Aitchison, A. Warren, D. Neal, P. Rabbitts, RASSF1A promoter methylation is frequently detected in both pre-malignant and nonmalignant microdissected prostatic epithelial tissues, *Prostate* 67 (2007) 638–644.
- [55] A. Flori, C. Steinhoff, M. M&rdquo;uller, H. Seifert, C. Hader, R. Engers, R. Ackermann, W. Schulz, Coordinate hypermethylation at specific genes in prostate carcinoma precedes LINE-1 hypomethylation, *Br. J. Cancer* 91 (2004) 985–994.
- [56] C. Goessl, H. Krause, M. M&rdquo;uller, R. Heicappell, M. Schrader, J. Sachsinger, K. Miller, Fluorescent methylation-specific polymerase chain reaction for DNA-based detection of prostate cancer in bodily fluids, *Cancer Res.* 60 (2000) 5941–5945.
- [57] P. Cairns, M. Esteller, J.G. Herman, M. Schoenberg, C. Jeronimo, M. Sanchez-Cespedes, N.-H. Chow, M. Grasso, L. Wu, W.B. Westra, et al., Molecular detection of prostate cancer in urine by GSTP1 hypermethylation, *Clin. Cancer Res.* 7 (2001) 2727–2730.
- [58] T. Jatkoe, R. Karnes, S. Freedland, Y. Wang, A. Le, J. Baden, A urine-based methylation signature for risk stratification within low-risk prostate cancer, *Br. J. Cancer* 112 (2015) 802–808.
- [59] T. Wu, E. Giovannucci, J. Welge, P. Mallick, W. Tang, S. Ho, Measurement of GSTP1 promoter methylation in body fluids may complement PSA screening: a meta-analysis, *Br. J. Cancer* 105 (2011) 65–73.
- [60] C. Haldrup, K. Mundbjerg, E.M. Vestergaard, P. Lamy, P. Wild, W.A. Schulz, C. Arsov, T. Visakorpi, M. Borre, S. Hoyer, et al., DNA methylation signatures for prediction of biochemical recurrence after radical prostatectomy of clinically localized prostate cancer, *J. Clin. Oncol.* 31 (2013) 3250–3258.
- [61] L. Richiardi, V. Fiano, C. Grasso, D. Zugna, L. Delsedime, A. Gillio-Tos, F. Merletti, Methylation of APC and GSTP1 in non-neoplastic tissue adjacent to prostate tumour and mortality from prostate cancer, *PLoS ONE* 8 (2013) 1–7.
- [62] R. Zelic, V. Fiano, D. Zugna, C. Grasso, L. Delsedime, L. Daniele, D. Galliano, A. Pettersson, A. Gillio-Tos, F. Merletti, et al., Global hypomethylation (LINE-1) and gene-specific hypermethylation (GSTP1) on initial negative prostate biopsy as markers of prostate cancer on a rebiopsy, *Clin. Cancer Res.* 22 (2015) 984–992.
- [63] N. Pertega-Gomes, J.R. Vizcaino, S. Felisbino, A.Y. Warren, G. Shaw, J. Kay, H. Whitaker, A.G. Lynch, L. Fryer, D.E. Neal, et al., Epigenetic and oncogenic regulation of SLC16A7 (MCT2) results in protein over-expression, impacting on signalling and cellular phenotypes in prostate cancer, *Oncotarget* 6 (2015) 21675–21684.
- [64] E. Sebestyén, M. Zawisza, E. Eyra, Detection of recurrent alternative splicing switches in tumor samples reveals novel signatures of cancer, *Nucleic acids Res.* 43 (2015) 1345–1356.
- [65] C.P. Nelson, R.K. La Creis, J. Sauvageot, W.B. Isaacs, A.M. De Marzo, J.D. Groopman, W.G. Nelson, T.W. Kensler, Protection against 2-hydroxyamino-1-methyl-6-phenylimidazo[4,5-b]pyridine cytotoxicity and DNA adduct formation in human prostate by glutathione S-transferase p1, *Cancer Res.* 61 (2001) 103–109.
- [66] M. Nakayama, C.J. Bennett, J.L. Hicks, J.I. Epstein, E.A. Platz, W.G. Nelson, A.M. De Marzo, Hypermethylation of the human glutathione S-transferase- $\pi$  gene (GSTP1) CpG island is present in a subset of proliferative inflammatory atrophy lesions but not in normal or hyperplastic epithelium of the prostate: a detailed study using laser-capture microdissection, *Am. J. Pathol.* 163 (2003) 923–933.
- [67] O.Y. Mian, M.H. Khatib, M. Hedayati, J. Coulter, B. Abubaker-Sharif, J.M. Schwaninger, R.K. Veeraswamy, J.D. Brooks, L. Hopkins, D.B. Shinohara, B. Cornblatt, W. Nelson, S. Yegnasubramanian, T. DeWeese, GSTP1 loss results in accumulation of oxidative DNA base damage and promotes prostate cancer cell survival following exposure to protracted oxidative stress, *Prostate* 76 (2016) 199–206.
- [68] R. Kanwal, M. Pandey, N. Bhaskaran, G.T. MacLennan, P. Fu, L.E. Ponsky, S. Gupta, Protection against oxidative dna damage and stress in human prostate by glutathione s-transferase p1, *Mol. Carcinog.* 53 (2014) 8–18.
- [69] L.B. Alexandrov, S. Nik-Zainal, D.C. Wedge, S.A. Aparicio, S. Behjati, A.V. Biankin, G.R. Bignell, N. Bolli, A. Borg, A.-L. Børresen-Dale, et al., Signatures of mutational processes in human cancer, *Nature* 500 (2013) 415–421.
- [70] J.M. Mosquera, H. Beltran, K. Park, T.Y. MacDonald, B.D. Robinson, S.T. Tagawa, S. Perner, T.A. Bismar, A. Erbersdobler, R. Dhir, et al., Concurrent AURKA and MYCN gene amplifications are harbingers of lethal treatment-related neuroendocrine prostate cancer, *Neoplasia* 15 (2013) 1–10.
- [71] H.-L. Tan, A. Sood, H.A. Rahimi, W. Wang, N. Gupta, J. Hicks, S. Mosier, C.D. Gocke, J.I. Epstein, G.J. Netto, et al., Rb loss is characteristic of prostatic small cell neuroendocrine carcinoma, *Clin. Cancer Res.* 20 (2014) 890–903.
- [72] H. Kinoshita, Y. Shi, C. Sandefur, L.F. Meisner, C. Chang, A. Choon, C.R. Reznikoff, G.S. Bova, A. Friedl, D.F. Jarrard, Methylation of the androgen receptor minimal promoter silences transcription in human prostate cancer, *Cancer Res.* 60 (2000) 3623–3630.
- [73] T. Nakayama, M. Watanabe, H. Suzuki, M. Toyota, N. Sekita, Y. Hirokawa, A. Mizokami, H. Ito, R. Yatani, T. Shiraishi, Epigenetic regulation of androgen

- receptor gene expression in human prostate cancers, *Lab. Invest.* 80 (2000) 1789–1796.
- [74] D.F. Jarrard, H. Kinoshita, Y. Shi, C. Sandefur, D. Hoff, L.F. Meisner, C. Chang, J.G. Herman, W.B. Isaacs, N. Nassif, Methylation of the androgen receptor promoter CpG island is associated with loss of androgen receptor expression in prostate cancer cells, *Cancer Res.* 58 (1998) 5310–5314.
- [75] G.L. Gravina, F. Marampon, M. Di Staso, P. Bonfili, A. Vitturini, E.A. Jannini, R.G. Pestell, V. Tombolini, C. Festuccia, 5-Azacytidine restores and amplifies the bicalutamide response on preclinical models of androgen receptor expressing or deficient prostate tumors, *Prostate* 70 (2010) 1166–1178.
- [76] G.L. Gravina, C. Festuccia, D. Millimaggi, V. Dolo, V. Tombolini, M. de Vito, C. Vicentini, M. Bologna, Chronic azacytidine treatment results in differentiating effects, sensitizes against bicalutamide in androgen-independent prostate cancer cells, *Prostate* 68 (2008) 793–801.
- [77] C.S. Zorn, K.J. Wojno, M.T. McCabe, R. Kuefer, J.E. Gschwend, M.L. Day, 5-aza-2-deoxycytidine delays androgen-independent disease and improves survival in the transgenic adenocarcinoma of the mouse prostate mouse model of prostate cancer, *Clin. Cancer Res.* 13 (2007) 2136–2143.
- [78] L.-C. Li, R. Chui, K. Nakajima, B.R. Oh, H.C. Au, R. Dahiya, Frequent methylation of estrogen receptor in prostate cancer: correlation with tumor progression, *Cancer Res.* 60 (2000) 702–706.
- [79] P.A. Jones, M.J. Wolkowicz, W.M. Rideout, F.A. Gonzales, C.M. Marziasz, G.A. Coetzee, S.J. Tapscott, De novo methylation of the MyoD1 CpG island during the establishment of immortal cell lines, *Proc. Natl. Acad. Sci. U.S.A.* 87 (1990) 6117–6121.
- [80] F. Antequera, J. Boyes, A. Bird, High levels of de novo methylation and altered chromatin structure at CpG islands in cell lines, *Cell* 62 (1990) 503–514.
- [81] D.J. Smiraglia, L.J. Rush, M.C. Fr&rdquo;uhwald, Z. Dai, W.A. Held, J.F. Costello, J.C. Lang, C. Eng, B. Li, F.A. Wright, M.A. Caligiuri, C. Plass, Excessive CpG island hypermethylation in cancer cell lines versus primary human malignancies, *Hum. Mol. Genet.* 10 (2001) 1413–1419.
- [82] J. Kawai, K. Hirose, S. Fushiki, Comparison of DNA methylation patterns among mouse cell lines by restriction landmark genomic scanning, *Mol. Cell. Biol.* 14 (1994) 7421–7427.
- [83] P.M. Vertino, J.P. Issa, O.M. Pereira-Smith, S.B. Baylin, Stabilization of DNA methyltransferase levels and CpG island hypermethylation precede SV40-induced immortalization of human fibroblasts, *Cell Growth Differ.* 5 (1994) 1395–1402.
- [84] S. Tommasi, A. Zheng, A. Wening, S.E. Bates, X.A. Li, X. Wu, M. Hollstein, A. Besaratinia, Mammalian cells acquire epigenetic hallmarks of human cancer during immortalization, *Nucleic Acids Res.* 41 (2013) 182–195.
- [85] S. Degerman, M. Landfors, J.K. Siwicki, J. Revie, M. Borss&eacute;n, E. Evel&rdquo;onn, E. Forestier, K.H. Chrzanoska, P. Ryd&eacute;n, W.N. Keith, G. Roos, Immortalization of T-cells is accompanied by gradual changes in CpG methylation resulting in a profile resembling a subset of T-cell leukemias, *Neoplasia* 16 (2014) 606–615.
- [86] K. Gordon, T. Clouaire, X.X. Bao, S.E. Kemp, M. Xenophontos, J.I. de Las Heras, I. Stancheva, Immortality, but not oncogenic transformation, of primary human cells leads to epigenetic reprogramming of DNA methylation and gene expression, *Nucleic Acids Res.* 42 (2014) 3529–3541.
- [87] K.E. Varley, J. Gertz, K.M. Bowling, S.L. Parker, T.E. Reddy, F. Pauli-Behn, M.K. Cross, B.A. Williams, J.A. Stamatoyannopoulos, G.E. Crawford, D.M. Absher, B.J. Wold, R.M. Myers, Dynamic DNA methylation across diverse human cell lines and tissues, *Genome Res.* 23 (2013) 555–567.
- [88] C.R. Jeter, B. Liu, X. Liu, X. Chen, C. Liu, T. Calhoun-Davis, J. Repass, H. Zaehres, J.J. Shen, D.G. Tang, NANOG promotes cancer stem cell characteristics and prostate cancer resistance to androgen deprivation, *Oncogene* 30 (2011) 3833–3845.
- [89] S. Kregel, R.Z. Szmulewitz, D.J. Vander Griend, The pluripotency factor NANOG is directly upregulated by the androgen receptor in prostate cancer cells, *Prostate* 74 (2014) 1530–1543.
- [90] T. Ueki, K.M. Walter, H. Skinner, E. J&eacute;e, R.H. Hruban, M. Goggins, Aberrant CpG island methylation in cancer cell lines arises in the primary cancers from which they were derived, *Oncogene* 21 (2002) 2114–2117.
- [91] M.F. Paz, M.F. Fraga, S. Avila, M. Guo, M. Pollan, J.G. Herman, M. Esteller, A systematic profile of DNA methylation in human cancer cell lines, *Cancer Res.* 63 (2003) 1114–1121.
- [92] P.T. Hennessey, M.F. Ochs, W.W. Mydlarz, W. Hsueh, L. Cope, W. Yu, J.A. Califano, Promoter methylation in head and neck squamous cell carcinoma cell lines is significantly different than methylation in primary tumors and xenografts, *PLoS ONE* 6 (2011) e20584.
- [93] C.E. Nestor, R. Ottaviano, D. Reinhardt, H.A. Cruickshanks, H.K. Mjoseng, R.C. McPherson, A. Lentini, J.P. Thomson, D.S. Dunican, S. Pennings, S.M. Anderson, M. Benson, R.R. Meehan, Rapid reprogramming of epigenetic and transcriptional profiles in mammalian culture systems, *Genome Biol.* 16 (2015) 11.
- [94] D. Cunningham, Z. You, In vitro and in vivo model systems used in prostate cancer research, *J. Biol. Methods* 2 (2015) 17.
- [95] A. Cock-Rada, J.B. Weitzman, The methylation landscape of tumour metastasis, *Biol. Cell* 105 (2013) 73–90.
- [96] M. Reynold, S. Turcan, D. Giri, K. Kannan, L.A. Walsh, A. Viale, M. Drobnjak, L.T. Vahdat, W. Lee, T.A. Chan, Remodeling of the methylation landscape in breast cancer metastasis, *PLoS ONE* 9 (2014) 1–10.
- [97] Y.-J. Kim, H.-Y. Yoon, S.-K. Kim, Y.-W. Kim, E.-J. Kim, I.Y. Kim, W.-J. Kim, EFEMP1 as a novel DNA methylation marker for prostate cancer: array-based DNA methylation and expression profiling, *Clin. Cancer Res.* 17 (2011) 4523–4530.
- [98] A.L. Statham, M.D. Robinson, J.Z. Song, M.W. Coolen, C. Stirzaker, S.J. Clark, Bisulfite sequencing of chromatin immunoprecipitated DNA (BisChIP-seq) directly informs methylation status of histone-modified DNA, *Genome Res.* 22 (2012) 1120–1127.
- [99] H. Takeshima, M. Wakabayashi, N. Hattori, S. Yamashita, T. Ushijima, Identification of coexistence of DNA methylation and H3K27me3 specifically in cancer cells as a promising target for epigenetic therapy, *Carcinogenesis* 36 (2015) 192–201.
- [100] P.C. Taberlay, A.L. Statham, T.K. Kelly, S.J. Clark, P.A. Jones, Reconfiguration of nucleosome-depleted regions at distal regulatory elements accompanies DNA methylation of enhancers and insulators in cancer, *Genome Res.* 2014, pp. 1421–1432.
- [101] V. Nogales, W.C. Reinhold, S. Varma, A. Martinez-Cardus, C. Moutinho, S. Moran, H. Heyn, A. Seb&eacute;o, A. Barnadas, Y. Pommier, M. Esteller, Epigenetic inactivation of the putative DNA/RNA helicase SLFN11 in human cancer confers resistance to platinum drugs, *Oncotarget* (2015).
- [102] C.P. Wong, A. Hsu, A. Buchanan, Z. Palomera-Sanchez, L.M. Beaver, E.A. Houseman, D.E. Williams, R.H. Dashwood, E. Ho, Effects of sulforaphane and 3, 3'-diindolylmethane on genome-wide promoter methylation in normal prostate epithelial cells and prostate cancer cells, *PLOS ONE* 9 (2014) e86787.
- [103] C.J. Phillip, C.K. Giardina, B. Bilir, D.J. Cutler, Y.H. Lai, O. Kucuk, C.S. Moreno, Genistein cooperates with the histone deacetylase inhibitor vorinostat to induce cell death in prostate cancer cells, *BMC Cancer* 12 (2012) 145.
- [104] N. Shukeir, B. Stefanska, S. Parashar, F. Chik, A. Arakelian, M. Szyf, S.A. Rabbani, Pharmacological methyl group donors block skeletal metastasis in vitro and in vivo, *Br. J. Pharmacol.* 172 (2015) 2769–2781.
- [105] P. Pakneshan, R.H. Xing, S.A. Rabbani, Methylation status of uPA promoter as a molecular mechanism regulating prostate cancer invasion and growth in vitro and in vivo, *FASEB J.* 17 (2003) 1081–1088.
- [106] L.C. Costello, R.B. Franklin, The clinical relevance of the metabolism of prostate cancer; zinc and tumor suppression: connecting the dots, *Mol. Cancer* 5 (2006) 17.
- [107] N.J. Serkova, E.J. Gamito, R.H. Jones, C. O'Donnell, J.L. Brown, S. Green, H. Sullivan, T. Hedlund, E.D. Crawford, The metabolites citrate, myo-inositol, and spermine are potential age-independent markers of prostate cancer in human expressed prostatic secretions, *Prostate* 68 (2008) 620–628.
- [108] G.F. Giske&oslash;deg&aring;rd, H. Bertilsson, K.M. Seln&aring;s, A.J. Wright, T.F. Bathen, T. Viset, J. Halgunset, A. Angelsen, I.S. Gribbestad, M.-B. Tessem, Spermine and citrate as metabolic biomarkers for assessing prostate cancer aggressiveness, *PLOS ONE* 8 (2013) e62375.
- [109] M. Clyne, Prostate cancer: biopsy citrate concentration could predict prostate cancer growth rate, *Nat. Rev. Urol.* 9 (2012), 123–123.
- [110] K.K. Singh, M.M. Desouki, R.B. Franklin, L.C. Costello, Mitochondrial aconitase and citrate metabolism in malignant and nonmalignant human prostate tissues, *Mol. Cancer* 5 (2006) 14.
- [111] H. Jakobsen, H. Rui, Y. Thomassen, T. Hald, K. Purvis, Polyamines and other accessory sex gland secretions in human seminal plasma 8 years after vasectomy, *J. Reprod. Fertil.* 87 (1989) 39–45.
- [112] H.E. Fj&oslash;se, H. Strand, A. Sunde, Dose-dependent induction of or nithine decarboxylase and S-adenosyl-methionine decarboxylase activity by testosterone in the accessory sex organs of male rats, *Prostate* 21 (1992) 239–245.
- [113] P.J. Blackshear, J.M. Manzella, D.J. Stumpo, L. Wen, J.K. Huang, O. Oyen, W.S. Young, High level, cell-specific expression of ornithine decarboxylase transcripts in rat genitourinary tissues, *Mol. Endocrinol.* 3 (1989) 68–78.
- [114] J. Cyriac, R. Haleem, X. Cai, Z. Wang, Androgen regulation of spermidine synthase expression in the rat prostate, *Prostate* 50 (2002) 252–261.
- [115] M. van der Graaf, R.G. Schipper, G.O. Oosterhof, J.A. Schalken, A.A. Verhofs-tad, A. Heerschap, Proton MR spectroscopy of prostatic tissue focused on the detection of spermine, a possible biomarker of malignant behavior in prostate cancer, *MAGMA* 10 (2000) 153–159.
- [116] R.C. Smith, M.S. Litwin, Y. Lu, B.R. Zetter, Identification of an endogenous inhibitor of prostatic carcinoma cell growth, *Nat. Med.* 1 (1995) 1040–1045.
- [117] W.A.M. Loenen, S-adenosylmethionine: jack of all trades and master of everything? *Biochem. Soc. Trans.* 34 (2006) 330–333.
- [118] S. Ottaviani, G.N. Brooke, C. O'Hanlon-Brown, J. Waxman, S. Ali, L. Buluwela, Characterisation of the androgen regulation of glycine N-methyltransferase in prostate cancer cells, *J. Mol. Endocrinol.* 51 (2013) 301–312.
- [119] A.P. Khan, T.M. Rajendiran, B. Ateeq, I.A. Asangani, J.N. Athanikar, A.K. Yocum, R. Mehra, J. Siddiqui, G. Palapattu, J.T. Wei, G. Michailidis, A. Sreekumar, A.M. Chinnaiyan, The role of sarcosine metabolism in prostate cancer progression, *Neoplasia* 15 (2013) 491–501.
- [120] A. Sreekumar, L.M. Poisson, T.M. Rajendiran, A.P. Khan, Q. Cao, J. Yu, B. Laxman, R. Mehra, R.J. Lonigro, Y. Li, M.K. Nyati, A. Ahsan, S. Kalyana-Sundaram, B. Han, X. Cao, J. Byun, G.S. Omenn, D. Ghosh, S. Pennathur, D.C. Alexander, A. Berger, J.R. Shuster, J.T. Wei, S. Varambally, C. Beecher, A.M. Chinnaiyan, Metabolomic profiles delineate potential role for sarcosine in prostate cancer progression, *Nature* 457 (2009) 910–914.
- [121] L. Sun, L. Song, Q. Wan, G. Wu, X. Li, Y.Y. Wang, J. Wang, Z. Liu, X. Zhong, X. He, S. Shen, X. Pan, A. Li, Y.Y. Wang, P. Gao, H. Tang, H. Zhang, cMyc-mediated activation of serine biosynthesis pathway is critical for cancer progression under nutrient deprivation conditions, *Cell Res.* 25 (2015) 429–444.
- [122] P.M. Tedeschi, E.K. Markert, M. Gounder, H. Lin, D. Dvorzhinski, S.C. Dolfi, L.L.-Y. Chan, J. Qiu, R.S. DiPaola, K.M. Hirshfield, L.G. Boros, J.R. Bertino, Z.N. Oltvai, A. Vazquez, Contribution of serine, folate and glycine metabolism to the ATP, NADPH and purine requirements of cancer cells, *Cell Death Dis.* 4 (2013) e877.

- [123] J. Cuzick, D.M. Berney, G. Fisher, D. Mesher, H. Møller, J.E. Reid, M. Perry, J. Park, A. Younus, A. Gutin, C.S. Foster, P. Scardino, J.S. Lanchbury, S. Stone, Prognostic value of a cell cycle progression signature for prostate cancer death in a conservatively managed needle biopsy cohort, *Br. J. Cancer* 106 (2012) 1095–1099.
- [124] D.C. Wallace, Mitochondria and cancer, *Nat. Rev. Cancer* 12 (2012) 685–698.
- [125] T.C. Larman, S.R. DePalma, A.G. Hadjipanayis, A. Protopopov, J. Zhang, S.B. Gabriel, L. Chin, C.E. Seidman, R. Kucherlapati, J.G. Seidman, Spectrum of somatic mitochondrial mutations in five cancers, *Proc. Natl. Acad. Sci. U.S.A.* 109 (2012) 14087–14091.
- [126] Y.S. Ju, L.B. Alexandrov, M. Gerstung, I. Martincorena, S. Nik-Zainal, M. Ramakrishna, H.R. Davies, E. Papaemmanuil, G. Gundem, A. Shlien, et al., Origins and functional consequences of somatic mitochondrial DNA mutations in human cancer, *Elife* 3 (2014).
- [127] R.S. Arnold, C.Q. Sun, J.C. Richards, G. Grigoriev, I.M. Coleman, P.S. Nelson, C.-L. Hsieh, J.K. Lee, Z. Xu, A. Rogatko, A.O. Osunkoya, M. Zayzafoon, L. Chung, J.A. Petros, Mitochondrial DNA mutation stimulates prostate cancer growth in bone stromal environment, *Prostate* 69 (2009) 1–11.
- [128] M.M. Nass, Differential methylation of mitochondrial and nuclear DNA in cultured mouse, hamster and virus-transformed hamster cells. In vivo and in vitro methylation, *J. Mol. Biol.* 80 (1973) 155–175.
- [129] B. Vanyushin, M. Kirnos, The nucleotide composition and pyrimidine clusters in DNA from beef heart mitochondria, *FEBS Lett.* 39 (1974) 195–199.
- [130] I.B. Dawid, 5-Methylcytidylic acid: absence from mitochondrial DNA of frogs and HeLa cells, *Science* 184 (1974) 80–81.
- [131] Y. Pollack, J. Kasir, R. Shemer, S. Metzger, M. Szyf, Methylation pattern of mouse mitochondrial DNA, *Nucleic Acids Res.* 12 (1984) 4811–4824.
- [132] L.R. Cardon, C. Burge, D.A. Clayton, S. Karlin, L.O.N.R. Cardont, C. Burget, D.A. Claytont, S. Karlint, Pervasive CpG suppression in animal mitochondrial genomes, *Proc. Natl. Acad. Sci. U.S.A.* 91 (1994) 3799–3803.
- [133] R.J. Shmookler Reis, S. Goldstein, Mitochondrial DNA in mortal and immortal human cells. Genome number, integrity, and methylation, *J. Biol. Chem.* 258 (1983) 9078–9085.
- [134] G.S. Groot, A.M. Kroon, Mitochondrial DNA from various organisms does not contain internally methylated cytosine in -CCGG- sequences, *Biochim. Biophys. Acta* 564 (1979) 355–357.
- [135] G. Agrimi, M.A. Di Noia, C.M. Marobbio, G. Fiermonte, F.M. Lasorsa, F. Palmieri, Identification of the human mitochondrial Sadenosylmethionine transporter: bacterial expression, reconstitution, functional characterization and tissue distribution, *Biochem. J.* 379 (2004) 183–190.
- [136] L.S. Shock, P.V. Thakkar, E.J. Peterson, R.G. Moran, S.M. Taylor, DNA methyltransferase 1, cytosine methylation, and cytosine hydroxymethylation in mammalian mitochondria, *Proc. Natl. Acad. Sci. U.S.A.* 108 (2011) 3630–3635.
- [137] H. Manev, S. Dzitoieva, H. Chen, Mitochondrial DNA: a blind spot in neuroepigenetics, *Biomol. Concepts* 3 (2012) 997–1003.
- [138] H. Chen, S. Dzitoieva, H. Manev, Effect of valproic acid on mitochondrial epigenetics, *Eur. J. Pharmacol.* 690 (2012) 51–59.
- [139] D. Bellizzi, P. DAquila, M. Giordano, A. Montesanto, G. Passarino, Global DNA methylation levels are modulated by mitochondrial DNA variants, *Epigenomics* 4 (2012) 17–27.
- [140] A. Maresca, DNA methyltransferase 1 mutations and mitochondrial pathology: is mtDNA methylated? *Front. Genet.* 6 (2015) 1–15.
- [141] A.A. Baccarelli, H.-M. Byun, Platelet mitochondrial DNA methylation: a potential new marker of cardiovascular disease, *Clin. Epigenet.* 7 (2015) 44.
- [142] S. Ghosh, S. Sengupta, V. Scaria, Comparative analysis of human mitochondrial methylomes shows distinct patterns of epigenetic regulation in mitochondria, *Mitochondrion* 18 (2014) 58–62.
- [143] M. Takasugi, S. Yagi, K. Hirabayashi, K. Shiota, DNA methylation status of nuclear-encoded mitochondrial genes underlies the tissue-dependent mitochondrial functions, *BMC Genomics* 11 (2010) 481.
- [144] P.F. Chinnery, H.R. Elliott, G. Hudson, D.C. Samuels, C.L. Relton, Epigenetics, epidemiology and mitochondrial DNA diseases, *Int. J. Epidemiol.* 41 (2012) 177–187.
- [145] R.D.W. Kelly, A. Mahmud, M. McKenzie, I.A. Trounce, J.C. St John, Mitochondrial DNA copy number is regulated in a tissue specific manner by DNA methylation of the nuclear-encoded DNA polymerase gamma A, *Nucleic Acids Res.* 40 (2012) 10124–10138.
- [146] W. Lee, J. Johnson, D.J. Gough, J. Donoghue, G.L.M. Cagnone, V. Vaghjiani, K.A. Brown, T.G. Johns, J.C. St. John, Mitochondrial DNA copy number is regulated by DNA methylation and demethylation of POLGA in stem and cancer cells and their differentiated progeny, *Cell Death Dis.* 6 (2015) e1664.
- [147] D.J. Smiraglia, M. Kulawiec, G.L. Bistulfi, S.G. Gupta, K.K. Singh, A novel role for mitochondria in regulating epigenetic modification in the nucleus, *Cancer Biol. Ther.* 7 (2008) 1182–1190.
- [148] S.R. Atilano, D. Malik, M. Chwa, J. Cáceres-Del-Carpio, A.B. Nesburn, D.S. Boyer, B.D. Kuppermann, S.M. Jazwinski, M.V. Miceli, D.C. Wallace, N. Udar, M.C. Kenney, Mitochondrial DNA variants can mediate methylation status of inflammation, angiogenesis and signaling genes, *Hum. Mol. Genet.* 24 (2015) 4491–4503.